

NITRO MATIC

17-5-11


GUARANTEED ANALYSIS:	By Wt.
TOTAL NITROGEN (N)*	17.00%
1.95% Ammoniacal Nitrogen	
5.57% Water Soluble Organic Nitrogen	
9.48% Water Insoluble Nitrogen	
AVAILABLE PHOSPHATE (P ₂ O ₅)	5.00%
AVAILABLE POTASH (K ₂ O)	11.00%
Boron (B)	0.05%
Total Copper (Cu)	0.05%
0.00% Soluble Copper (Cu)	
Chlorine (Cl), not more than	2.00%
Total Iron (Fe)	1.00%
0.7% Soluble Iron (Fe)	
0.1% Chelated Iron (Fe)	
Total Magnesium (Mg)	1.50%
1.50% Soluble Magnesium (Mg)	
Total Manganese (Mn)	0.50%
0.39% Soluble Manganese (Mn)	
Molybdate (Mo)	0.01%
Total Zinc (Zn)	0.12%
0.12% Soluble Zinc (Zn)	

DERIVED FROM: Ureaform, Ammonium Phosphate, Sulfate of Potash, Sulfate of Potash Magnesia, Borax, Copper Sulfate, Iron EDTA/ Iron DPTA, Iron Sulfate, Manganese Sulfate, Manganese Sulfate, Sodium Molybdate and Zinc Sulfate.

GUARANTEED BY: Southern Agricultural Insecticides, Inc. P.O. Box 218 Palmetto, FL 34220

12 Units of slow release nitrogen from NITROFORM™

**CAUTION: KEEP OUT OF REACH OF CHILDREN.
MAY BE HARMFUL IF SWALLOWED. AVOID CONTACT WITH EYES.**



Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28793

Boone, NC 28607

F 672

Net Weight: 50 lbs.(22.68 kg.)

BLUE CHIP

NURSERY SPECIAL

NURSERY

CONTAINER TOPDRESS RATES:

1 gallon *	1 level or heaping Tablespoon
3 gallon	2 Tablespoons (level 1/8 cup)
5- 7 gallon	heaping 1/8 cup
10 gallon	level 1/4 cup

* Use on established 1 gallon crops only.

Starter Incorporation: 6- 8 pounds per cubic yard.

NOTE: Rates depend upon crop, irrigation frequency, temperature and total root mass. Never apply fertilizer near the stems. It is best to make a ring of fertilizer in the outer root zone. Adjust soil pH prior to potting if necessary. If plants are being started from seed, have low root mass, or are immature when planted, use 1/2 the rates given above.

LANDSCAPE

ORNAMENTALS: On established shrubs, woody ornamentals, small trees, and ground covers in the landscape, apply at approximately 1 cup per 30 sq. ft.

TREES: On established trees in the ground: apply 1/4 cup per tree for root balls equal to 24 inches, apply 1/2 cup for root balls equal to 36 inches, or for larger trees use 1/4 - 1/2 cup per inch of tree diameter.

For broadcasting along rows: apply at 30 lbs. per 1,000 sq. ft.

LAWN: Great for recently sprigged or plugged lawns!

Apply with spreader at 5 - 5.5 lbs. per 1,000 sq. ft.

FOR FLORIDA:

We recommend that you follow the Green Industries BMP's at:

http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/BMP_Book_final.pdf

Fertilizers containing secondary mineral elements may stain masonry surfaces. After applying, sweep fertilizer from sidewalks and other masonry surfaces.

Timing, method and rates of application, differences in plant varieties, growing conditions and other influencing factors are beyond the control of the seller.

Buyer assumes all risks of use, storage or handling of this material.