

THIS PRODUCT CONTAINS:

0.258% 2,4-dichlorophenoxyacetic acid equivalent 0.062% (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.028% 3,6-dichloro-o-anisic acid equivalent
Isomer Specific by AOAC Methods.

CONTAINS SINGLE ISOMER FORM OF MECOPROP-p

2,4-D, dimethylamine salt (CAS Registry No. 2008-39-1)
Mecoprop-p, dimethylamine salt (CAS Registry No. 66423-09-4)
Dicamba, dimethylamine salt (CAS Registry No. 2300-66-5)

CAS Registry Numbers:
2,4-D, dimethylamine salt CAS# 2008-39-1
Mecoprop-p, dimethylamine salt CAS# 66423-09-4
Dicamba, dimethylamine salt CAS#2300-66-5

S A SOUTHERN 5 0 AG® LAWN SPOT WEEDER

READY-TO-USE

KEEP OUT OF REACH OF CHILDREN

CAUTION

See inside of label for additional precautions, First Aid and Directions for Use

ACTIVE INGREDIENTS:

2,4-D, dimethylamine salt 0.311%
Mecoprop-p, dimethylamine salt 0.075%
Dicamba, dimethylamine salt 0.034%
INERT INGREDIENTS:..... 99.580%
TOTAL 100.000%

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes or clothing. Wear goggles when applying this product. When using this product, wear long-sleeved shirt, long pants, socks, shoes and rubber gloves.

After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower.

Distributed by **Southern Agricultural Insecticides, Inc.**

Palmetto, FL 34220 Hendersonville, NC 28793 Boone, NC 28607
E.P.A. Reg. No. 2217-585-829 J1

Net Contents Liquid: 24 ozs. (709 ml)

EPA EST. NO. 829-FL-1 (3);
829-NC-1 (4); 829-NC-2 (5)
Number following Est. No.
corresponds to first digit in
Lot Number

First Aid

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep from freezing. Store in original container in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL: If Empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partially filled: Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRODUCT DESCRIPTION: Spot Weeder contains TRIMEC® turf herbicide, a combination of three proven weed killers - 2,4-D, mecoprop-p, and dicamba. Together they control a wide range of broadleaf weeds commonly found in home (residential) lawns. This ready-to-use herbicide is intended for spot treatments only. Weeds that have not emerged at the time of application will not be controlled.

WHERE TO USE: Northern Grasses - Kentucky bluegrass, perennial ryegrass and fescues. Southern Grasses - St. Augustinegrass, centipedegrass, bentgrass, bahiagrass, bermudagrass and zoysiagrass

PROHIBITIONS: Do not apply to garden plants, flowers or shrubs. Do not use this product on carpetgrass, dichondra, nor on lawns with desirable clovers.

HOW TO USE: Use to spot treat weeds in the lawn. Spot Weeder is designed for easy treatment of small weed patches without mess or trouble and without bending or stooping.

ENVIRONMENTAL HAZARDS:

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

NO MIXING OR POURING - AIM AT WEED AND SPRAY - For best results, use Spot Weeder when weeds are actively growing. Rainfall or watering soon after application may reduce effectiveness.

FOR CONTAINERS WITH TRIGGER SPRAYERS: Spot Weeder is ready-to-use and is applied directly from the container fitted with the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Usually, two (2) pumps of this trigger sprayer will provide adequate coverage of individual weeds. Overspraying may injure lawns including bentgrass, centipede grass, and St. Augustine grass.

FOR REFILL CONTAINERS: Spot Weeder is ready-to-use and is applied directly from the container. Remove the cap from the container and attach the trigger sprayer. Direct the spray to the target weeds to provide uniform coverage of the upper leaf surfaces. Overspraying may injure lawns including bentgrass, centipede grass, and St. Augustine grass.

CARE AND INSTRUCTIONS FOR TRIGGER SPRAYERS - OPERATION: 1 Remove sprayer from bag. 2. Unscrew cap from bottle and remove the foil seal. 3. Making sure tube is properly inserted through hole in cap, insert tube into bottle and screw on cap. 4. Open spray nozzle approximately one full turn. 5. TO PRIME - With sprayer directed at the target area, operate trigger until spray begins (approximately 10 to 15 times). 6. Adjust the sprayer nozzle for desired spray pattern.

STORAGE OF TRIGGER SPRAYER: 1. Close spray nozzle. 2. Hook the trigger on or under the bottle handle so that trigger is above the level of contents. 3. To prevent clogging of trigger, remove sprayer from bottle and pump clean water through the hose and trigger.

HOW TO PRIME AND CARE FOR YOUR SPRAYER:

1. After shaking product well, remove cap from container and replace with sprayer hose and cap assembly. Tighten securely. 2. Turn sprayer nozzle one half turn counter-clockwise. Hold sprayer below the level of container and prime by pumping trigger 10 - 15 times until product is dispensed. Adjust nozzle to coarse stream. 3. To store, turn nozzle to "OFF" position and store with sprayer above level of liquid in container to prevent leakage

Use Spot Weeder to kill poison ivy and any other troublesome broadleaf weeds.

- WEEDS -

Annual yellow sweetclover	Cockle	Musk thistle	Spiny cocklebur
Aster	Cocklebur	Mustard	Spiny sowthistle
Austrian fieldcress	Common mullein	Narrowleaf plantain	Spotted catsear
Bedstraw	Creeping jenny	Narrowleaf vetch	Spotted spurge

③

(WEEDS continued)

Carpentweed	Horsenettle	Russian thistle	Wild rape
Catchweed bedstraw	Jimsonweed	Scarlet pimpernel	Wild strawberry
Catnip	Knawel	Scotch thistle	Wild sweet potato
Catsear	Knotweed	Sheep sorrel	Wild vetch
Chickweed, common	Kochia	Shepherdspurse	Woodsorrel
Chickweed, mouseear	Lambsquarters	Slender plantain	Woolly croton
Chicory	Lespedeza	Smallflower galinsoga	Woolly morningglory
Cinquefoil	Mallow	Smooth dock	Woolly plantain
Clover, crimson	Matchweed	Smooth pigweed	Wormseed
Clover, hop	Mexicanweed	Sorrel	Yarrow
Clover, red	Milk vetch	Sowthistle	Yellow rocket
Clover, strawberry	Morningglory	Spanishneedles	Yellowflower pepperweed
Clover, sweet	Mouseear hawkweed	Speedwell	and other
Clover, white	Mugwort	Spiny amaranth	broadleaf weeds

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

For information and questions concerning this product please contact us ⑤ by email at support@southernag.com.

(WEEDS continued)

Beggarlicks	Cudweed	Nettle	Spurweed
Betony, Florida	Curly dock	Orange hawkweed	Stinging nettle
Bindweed, field	Daisy, English	Oriental cocklebur	Strawberry, India
mock	Bird vetch	Daisy fleabane	Oxalis Tall nettle
Bitter wintercress	Daisy, oxeye	Parsley-piert	Tall vervain
Bittercress, hairy	Dandelion	Parsnip	Tansy ragwort
Bitterweed	Dichondra	Pearlwort	Tansy mustard
Black-eyed Susan	Dogbane	Pennycress	Tanweed
Black medic	Dogfennel	Pennywort	Thistle
Black mustard	Dollarweed	Pepperweed	Trailing crownvetch
Blackseed plantain	Elderberry	Pigweed	Tumble mustard
Blessed thistle	False dandelion	Pineywoods bedstraw	Tumble pigweed
Bloodflower milkweed	Falseflax	Plains coreopsis (lickseed)	Velvetleaf
Blue lettuce	False sunflower	Plantain	Venice mallow
Blue vervain	Fiddleneck	Poison ivy	Virginia buttonweed
Bracted plantain	Florida pusley	Poison oak	Virginia creeper
Brassbuttons	Frenchweed	Pokeweed	Virginia pepperweed
Bristly oxtongue	Galinsoga	Poorjoe	Water pennywort
Broadleaf dock	Goathead	Prairie sunflower	Wavyleaf bullthistle
Broadleaf plantain	Goldenrod	Prickly lettuce	Western clematis
Broomweed	Ground ivy	Prickly sida	Western salsify
Buckhorn	Gumweed	Prostrate knotweed	White mustard
Buckhorn plantain	Hairy fleabane	Prostrate pigweed	Wild aster
Bulbous buttercup	Hawkweed	Prostrate spurge	Wild buckwheat
Bull thistle	Healall	Prostrate vervain	Wild carrot
Bullnettle	Heartleaf drymaria	Puncturevine	Wild four-o'clock
Burclover	Heath aster	Purslane, common	Wild garlic
Burdock	Hedge bindweed	Ragweed	Wild geranium
Burning nettle	Hedge mustard	Red sorrel	Wild lettuce
Burweed	Hemp	Redroot pigweed	Wild marigold
Buttercup	Henbit	Redstem filaree	Wild mustard
Buttonweed	Hoary cress	Rough cinquefoil	Wild onion
Canada thistle	Hoary plantain	Rough fleabane	Wild parsnip
Carolina geranium	Hoary vervain	Russian pigweed	Wild radish

④