

TOMATO DUST

FOR USE ONLY ON HOME VEGETABLE GARDENS.

ACTIVE INGREDIENT:	By Wt.
BASIC COPPER SULFATE*	13.30%
INERT INGREDIENTS:	86.70%
TOTAL:	100.00%
*Metallic Copper Equivalent	7.10%



KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Seek medical attention as soon as possible if irritation persists. **IF INHALED:** Remove victim to fresh air. Contact a physician if irritation occurs. **IF SWALLOWED:** Give victim 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention. **IF ON SKIN:** Wash affected area with plenty of soap and water. Contact a physician if irritation occurs.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: HARMFUL IF SWALLOWED. AVOID CONTACT WITH EYES, SKIN, OR CLOTHING. AVOID BREATHING DUST. WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING. MAY CAUSE SKIN SENSITIZATION REACTIONS IN CERTAIN INDIVIDUALS.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic organisms. Do not apply directly to water. Do not contaminate water by disposal of equipment washwaters. Drift and runoff from treated areas may be hazardous to fish and aquatic organisms in adjacent aquatic sites. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

A FUNGICIDE DUST FOR USE ON: TOMATOES and POTATOES for the control of early and late blights and as an aid in controlling bacterial spot. Dust plants thoroughly at weekly intervals and following rains. For potatoes make 11-22 ozs. per 1,000 sq. ft. applications beginning when plants are 6 inches high. For tomatoes make 11-22 ozs. per 1,000 sq. ft. applications beginning when plants are set in the field.

CUCUMBERS, SQUASH, WATERMELONS and CANTALOUPEs for the control of downy mildew. Dust thoroughly, but lightly, using 5 1/2 - 15 ozs. per 1,000 sq. ft. Apply at 10 day intervals and following rains. Heavy applications, applications to tender foliage, and applications to wet foliage can cause injury to cucurbits.

PEPPER as an aid in the control of bacterial spot, apply at weekly intervals and if possible preceding heavy rains, using 7 1/3 - 16 1/2 ozs. per 1,000 sq. ft. applications. There is no time limitation on the use of this dust. Remove excess residue at harvest. Consult proper State Extension specialist to determine if local recommendations are consistent with this label before using. Do not exceed indicated dosages. May cause injury to tender plant parts particularly when applications are excessive, uneven, or when foliage is wet. Control of the indicated diseases may not be satisfactory during periods of frequent or heavy rainfall, when disease infestations are extremely severe, or when applications are uneven, inadequate, or poorly timed.

Do not apply or allow drift to corn, grain sorghum, or other copper sensitive plants. Do not apply when conditions favor drift from treated area.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store pesticide in the original container in a locked storage area out of the reach of children and pets.

PESTICIDE DISPOSAL: To dispose of pesticide, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

Buyer assumes all risks of use, storage, or handling of this material in strict accordance with the directions and precautions given herewith.

EPA EST. No. 829-FL-1 (3); 829-NC-1 (4); 829-NC-2(5) Number following establishment No. corresponds to first digit in lot No.

E.P.A. REG. No. 829-6

Southern Agricultural Insecticides, Inc.

Palmetto, FL 34220

Hendersonville, NC 28793

Boone, NC 28607

revised 1998