



MOLE CRICKET & CHINCH BUG LAWN INSECT CONTROL

CONTAINS 0.1% DELTAMETHRIN FOR LAWN AND HOME INVADING PESTS

ACTIVE INGREDIENT:

Deltamethrin: 0.1%
OTHER INGREDIENTS..... 99.9%
100.0%

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use in commercial greenhouses, nurseries, on sod farms or grass grown for seed or plants being grown for sale. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

General Use Instructions

- 1. Determine size of area to be treated - measure and multiply length times width.
2. Determine appropriate spreader setting using chart.
3. Treat as directed - thorough coverage is important. Water in thoroughly with a half inch of water.
4. Keep children and pets off treated areas until this material is washed into turf and grass is dry.

OVERALL LAWN TREATMENT FOR GRASS AND DICHONDRA LAWNS

TO CONTROL Ants, Armyworms (inc. fall, Yellow-Striped, etc), Bermudagrass Mites, Pillbugs (adult), Boxelder Bugs, Centipedes, Chiggers, Chinch Bugs, Clover Mites, Cockroaches, Crickets, Cutworms, Darkling Ground Beetles, Digger Wasps, Essex Skipper, European Crane Fly Larvae, Field Crickets, Fiery Skipper, Fire Ants, Flea Beetle (in dichondra), Grasshopper, Hyperodes (Turfgrass Weevil), Japanese Beetle (adult), Lawn Moths, Leafhoppers, Adult Mealybugs, Silverfish & Firebrats, Sod Webworms, Sowbugs, Spiders, Spittlebugs, Springtails, Vegetable Weevils (in dichondra); Apply this product uniformly to home lawns, under shrubbery, and other ornamental recreational turf grass areas.

EUROPEAN CRANEFLY LARVAE (LEATHERJACKETS) - the best time to apply is in early April when the insects are young. Re-treat if infestation occurs, however do not retreat more than once every 2 weeks.

CHINCH BUGS - Start treatments when temperatures reach 80°F or above. A second application may be necessary in 6 weeks if insect activity continues. Re-treat if infestation occurs, however do not retreat more than once every 2 weeks.

SOD WEBWORMS (CRAMBUS spp.) - Sod Webworms are the light brown caterpillars 1/4 to 3/4 inch long, that hide in the grass during the day and feed on the grass blades at night. Adult Webworms, usually referred to as Lawn Moths (about 1 inch in length with long snout) emerge from the grass at dusk. They fly over the lawn laying eggs that later hatch into destructive caterpillars. Damage appears first as small brown spots which, if not treated, may rapidly develop into large brown areas of severely damaged or dead grass.

BAND (PERIMETER) TREATMENT AROUND HOUSE FOUNDATION: To help prevent infestation of buildings by Ants, Boxelder Bugs, brown dog ticks, Centipedes, Clover Mites, Cockroaches, Crickets, Field Crickets, Sowbugs, Spiders, Springtails (Collembola), and Ticks; treat a band 5 to 10 feet wide around and adjacent to the building where pests are active and may find entrance. Treat at a rate of 2 to 3 lb. per 1,000 sq. ft. Re-treat if infestation occurs, however do not retreat more than once every 2 weeks.

FLOWER, SHRUB AND ORNAMENTAL PLANT BEDS: To control Ants, Fire Ants, Cockroaches, Crickets, Cutworms, Sowbugs, Pillbugs, Grasshoppers, Armyworms, Brown Dog Ticks and Springtails apply granules evenly over soil within the beds and under shrubbery. Use the same dosage recommended for control of these pests in lawns. Water thoroughly with 1/2 inch of water. Re-treat if infestation occurs, however do not retreat more than once every 2 weeks.

SPOT TREATMENT ANTS NESTING OUTDOORS: This product may be used outdoors for the control of ants, including: Acrobat, Argentine, Big Headed, Cornfield, Field, Harvester, Leafcutter, Little Black, Pavement and Pyramid Ants. If possible, it is best to try to locate nest and treat this first. Nests will usually be found in the ground but may be concealed under stones, boards, and slabs or in stumps or logs. Product should be applied to thoroughly cover the nest (1 to 2 teaspoons) and sprinkled gently with sufficient water to dissolve granules.

IMPORTED FIRE ANTS: Apply two tablespoons over each 8' to 10' diameter mound and up to 2 feet around to prevent escape of the queen. Immediately after application, water the granules into the soil gently with a light sprinkling to avoid disturbing the colony. Use at least 1 gallon of water per mound, but no more than 2 gallons. Use of high pressure watering equipment may disturb the colony and cause migration of the ants, thus reducing the effectiveness of the treatment.

SPREADER SETTINGS

Note: Spreader settings given are approximate and variation can occur between identical models. User should calibrate spreader by accurately weighing a few pounds into your spreader and applying to a measured area. Then adjust spreader setting if necessary to assure correct rate of application.

Setting to deliver approximately: Spreader Make and Model

Table with 2 columns: Spreader Make and Model, 2 lb./ 1,000 sq. ft., 3 lb./ 1,000 sq. ft. Rows include Scotts Drop Spreaders, Scotts EasyGreen Rotary, Republic EZ Drop, Cyclone Model B1 Rotary, and Sears Craftsman Broadcast.

IMPORTANT. READ BEFORE USE

By using this product, user or buyer accepts the following conditions, disclaimer of warranties and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be adequate and should be followed carefully. However, because of manner of use and other factors beyond the control of Southern Agricultural Ins. (manufacturer), it is impossible for manufacturer to eliminate all risks associated with the use of this product.

In case of Medical emergencies or health and safety inquiries or in case of fire, leaking or damaged containers, information may be obtained by calling 1-800-471-0660.

DISCLAIMER OF WARRANTIES: THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Southern Agricultural Ins. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein.

LIMITATIONS OF LIABILITY: THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT MANUFACTURER'S ELECTION, THE REPLACEMENT OF PRODUCT.

For product information email us at sales@Southernag.com EPA EST. NO 829-FL-1 (3); 829-NC-1 (4); 829-NC-2 (5) Number following Est. No. Corresponds to first digit in Lot No.

Southern Agricultural Insecticides, Inc. Palmetto, Fla. 34220 Hendersonville, N.C. 28793 Boone, N.C. 28607 E.P.A. Reg. No. 829 - 294